



ELSEVIER Linear Algebra and its Applications 302–303 (1999) 295–302

**LINEAR ALGEBRA
AND ITS
APPLICATIONS**

www.elsevier.com/locate/laa

A characterization of trace zero nonnegative 5×5 matrices

Thomas J. Laffey^{*}, Eleanor Meehan

Department of Mathematics, University College Dublin, Belfield, Dublin, Ireland

Received 23 October 1998; accepted 22 May 1999

Submitted by M. Goldberg

Dedicated to Hans Schneider

Abstract

Let $\sigma = (\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5)$ be a list of complex numbers with $\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 + \lambda_5 = 0$. Necessary and sufficient conditions for the existence of an entry-wise nonnegative 5×5 matrix A with spectrum σ are presented. © 1999 Elsevier Science Inc. All rights reserved.

Keywords: Matrix; Spectrum; Graph; Cycles; Petersen graph; Nonnegative inverse eigenvalue problem

1. Introduction

Given a list $\sigma = (\lambda_1, \dots, \lambda_n)$ of complex numbers, the nonnegative inverse eigenvalue problem (NIEP) asks for the determination of necessary and sufficient conditions for the existence of an entrywise nonnegative $n \times n$ matrix A with spectrum σ . In this paper the problem is solved for $n = 5$ in the “trace zero” case, that is, when $\lambda_1 + \dots + \lambda_5 = 0$. Previous results on this problem for ‘small’ matrices include the case $n = 3$ (without further restriction) by Loewy and London [3], $n = 4$ and σ a list of real numbers also by Loewy and London [3], $n = 4$ and $\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 = 0$ by Reams [6]. Several other special cases of the NIEP for $n = 4$ have also been resolved by Reams [5]. For more results on the NIEP see [1,4].

^{*} Corresponding author. E-mail: laffey@ollamh.ucd.ie

2. An inequality

We now consider the NIEP for $n = 5$ when the sum of the five complex numbers is zero. So let $\sigma = (\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5)$ be a list of complex numbers. Define

$$s_k := \lambda_1^k + \lambda_2^k + \lambda_3^k + \lambda_4^k + \lambda_5^k, \quad k = 1, 2, 3, \dots$$

and assume $s_1 = 0$. We wish to determine necessary and sufficient conditions so that there exists a nonnegative 5×5 matrix A with spectrum $\sigma(A) = \sigma$. Suppose that such an A exists. Then,

$$s_k = \text{tr}(A^k) \quad \text{for } k = 1, 2, 3, \dots$$

Note that since A is entrywise nonnegative, $s_k \geq 0$ for all k . Since $s_1 = 0$, all the diagonal entries of A are zero. In this section we prove the following theorem.

Theorem 2.1. *Let A be an entrywise nonnegative 5×5 matrix with $\text{tr}(A) = 0$. Then,*

$$12s_5 - 5s_2s_3 + 5s_3\sqrt{4s_4 - s_2^2} \geq 0.$$

Proof. The result is clear if $12s_5 \geq 5s_2s_3$, since $4s_4 - s_2^2 \geq 0$ by [2], so we first analyse the circumstances in which $12s_5 < 5s_2s_3$. Let $A = (a_{ij})$. Write x_1, x_2, \dots, x_{10} for the 2-cycle products $a_{12}a_{21}, a_{13}a_{31}, \dots, a_{45}a_{54}$. So,

$$s_2 = 2(x_1 + x_2 + \dots + x_{10}).$$

There are $\binom{5}{3} = 10$ 3-subsets $\{\alpha, \beta, \gamma\}$ of $\{1, 2, 3, 4, 5\}$ and each 3-subset of symbols gives rise to two 3-cycles in s_3 . For example, $\{1, 2, 3\}$ gives the terms $a_{12}a_{23}a_{31}$ and $a_{13}a_{32}a_{21}$ and each occurs in s_3 with a multiplicity of 3. We can identify each 3-subset as the complement of the corresponding 2-subset. Denote these 3-cycle contributions by y_1, y_2, \dots, y_{10} where if $x_i = a_{rs}a_{sr}$, say, y_i corresponds to the 3-cycles from the complement of $\{r, s\}$ in $\{1, 2, 3, 4, 5\}$. For example, if $x_1 = a_{12}a_{21}$ then $y_1 = a_{34}a_{45}a_{53} + a_{35}a_{54}a_{43}$. Note that

$$s_3 = 3(y_1 + y_2 + \dots + y_{10}).$$

Thus,

$$5s_2s_3 = 30(x_1 + x_2 + \dots + x_{10})(y_1 + y_2 + \dots + y_{10}).$$

Next consider s_5 . Since A has a zero diagonal, s_5 is made up of contributions of the form (2-cycle) \times (3-cycle) and 5-cycles. For example, $a_{12}a_{21}a_{14}a_{43}a_{31}$ and $a_{12}a_{23}a_{34}a_{45}a_{51}$ are terms occurring in the $(1, 1)$ entry of A^5 . The general $(1, 1)$ entry of A^5 is $a_{1k}a_{kl}a_{lm}a_{mn}a_{n1}$ with $k \neq 1, k \neq l, l \neq m, m \neq n, n \neq 1$ (otherwise the

term is zero). In this position, if the term is not a 5-cycle type term, then either $l = 1$ or $m = 1$. Note that a term of the form $a_{12}a_{21}a_{34}a_{45}a_{53}$ cannot occur in s_5 . In terms of the x 's, and y 's previously defined, $x_1y_1, x_2y_2, \dots, x_{10}y_{10}$ do not occur in s_5 (since by our choice of notation, the 3-cycles involved in y_i are made from the symbols complementary to the symbols forming the 2-cycle in x_i). Hence in $12s_5 - 5s_2s_3$ we get the term $x_1y_1 + x_2y_2 + \dots + x_{10}y_{10}$ with coefficient -30 .

Consider now a product x_iy_j where $i \neq j$. Suppose that x_i corresponds to the 2-cycle term generated by the symbols $\{1, 2\}$, so y_j corresponds to the terms generated by the symbols $\{p, q, r\}$, where $\{p, q, r\}$ contains at least one symbol of $\{1, 2\}$.

Case 1: $p = 1, q = 2$. Say $r = 3$. In this case $y_j = a_{12}a_{23}a_{31} + a_{13}a_{32}a_{21}$. Note that $a_{12}a_{21}a_{12}a_{23}a_{31}$ and $a_{12}a_{21}a_{13}a_{32}a_{21}$ both occur in s_5 with coefficient 5 and thus these terms occur in $12s_5 - 5s_2s_3$ with coefficient $12 \times 5 - 30 = 30$.

Case 2: $\{1, 2\} \cap \{p, q, r\}$ contains one element. Suppose $p = 1$ is common and let $q = 3, r = 4$. We find that s_5 contains $a_{12}a_{21}a_{13}a_{34}a_{41}$ and $a_{12}a_{21}a_{14}a_{43}a_{31}$ both with coefficient 5. So these terms occur in $12s_5 - 5s_2s_3$ with coefficient $+30$.

The same arguments apply with symbols 1 and 2 replaced by p , and q , respectively. Hence,

$$12s_5 - 5s_2s_3 \geq 30 \sum_{i=1}^{10} \sum_{j \neq i, j=1}^{10} x_iy_j - 30 \sum_{i=1}^{10} x_iy_i$$

(\geq because we have excluded the 5-cycles).

We are supposing that $12s_5 - 5s_2s_3 < 0$. Then,

$$x_1y_1 + x_2y_2 + \dots + x_{10}y_{10} > \sum_{i=1}^{10} \sum_{j \neq i, j=1}^{10} x_iy_j. \quad (2.1)$$

Let x_1 be the maximum of x_1, x_2, \dots, x_{10} and suppose that

$$x_1 \leq x_2 + \dots + x_{10}.$$

Now the right-hand side of (2.1) contains $x_1(y_2 + \dots + y_{10}) + y_1(x_2 + \dots + x_{10})$. But,

$$\begin{aligned} x_1(y_2 + \dots + y_{10}) + y_1(x_2 + \dots + x_{10}) &\geq x_1(y_2 + \dots + y_{10}) + y_1x_1 \\ &\geq x_1y_1 + x_2y_2 + \dots + x_{10}y_{10} \end{aligned}$$

since $x_2 \leq x_1, \dots, x_{10} \leq x_1$. However, this says that the right-hand side of (2.1) is greater than or equal to the left-hand side which is false. Hence, we have proved.

Step 1: If $12s_5 < 5s_2s_3$ and x_1 is the maximum of x_1, x_2, \dots, x_{10} then

$$x_1 > x_2 + x_3 + \dots + x_{10}. \quad (2.2)$$

We now analyse $4s_4 - s_2^2$ under the assumption (2.2). Recall [2] that the part of $4s_4 - s_2^2$ which does not involve 4-cycles can be written as

$$4(x_1 \cdots x_{10})(J - 2P) \begin{pmatrix} x_1 \\ \vdots \\ x_{10} \end{pmatrix}$$

where J is the matrix of all “ones” and P is the adjacency matrix of the Petersen graph P_{10} . Note that we can label x_1, \dots, x_{10} so that P is

$$\begin{pmatrix} C + C^T & I \\ I & C^2 + (C^T)^2 \end{pmatrix}$$

where

$$C = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

Now

$$\begin{aligned} & (x_1 \cdots x_{10})(J - 2P) \begin{pmatrix} x_1 \\ \vdots \\ x_{10} \end{pmatrix} \\ &= (x_1 + \cdots + x_{10})^2 - 4[x_1x_2 + x_1x_5 + x_1x_6 + x_2x_3 + x_2x_7 + x_3x_4 \\ &\quad + x_3x_8 + x_4x_5 + x_4x_9 + x_5x_{10} + x_6x_8 + x_6x_9 + x_7x_9 + x_7x_{10} + x_8x_{10}] \\ &= (x_1 - (x_2 + \cdots + x_{10}))^2 + 4x_1 - 4[x_2x_3 + x_2x_7 + x_3x_4 + x_3x_8 + x_4x_5 \\ &\quad + x_4x_9 + x_5x_{10} + x_6x_8 + x_6x_9 + x_7x_9 + x_7x_{10} + x_8x_{10}]. \end{aligned}$$

Note that each negative term x_px_q has either p or q in $\{3,4,7,8,9,10\}$ and all the negative terms arise in the expansion of

$$(x_2 + x_3 + \cdots + x_{10})(x_3 + x_4 + x_7 + x_8 + x_9 + x_{10}).$$

Since

$$x_1 > x_2 + x_3 + \cdots + x_{10}$$

we get that $x_1(x_3 + x_4 + x_7 + x_8 + x_9 + x_{10}) \geq$ the sum of the terms x_px_q occurring with negative signs. Hence,

$$4s_4 - s_2^2 \geq 4(x_1 \cdots x_{10})(J - 2P) \begin{pmatrix} x_1 \\ \vdots \\ x_{10} \end{pmatrix} \geq 4[x_1 - (x_2 + \cdots + x_{10})]^2.$$

Hence we have established.

Step 2: If $x_1 > x_2 + \cdots + x_{10}$, then $4s_4 - s_2^2 \geq 4[x_1 - (x_2 + \cdots + x_{10})]^2$. (It is worth noting that the pairing off of the negative terms $x_p x_q$ with positive terms $x_r x_s$ with $s \in \{3, 4, 7, 8, 9, 10\}$ is crucial for the inequality. For example, if say $x_5 x_6$ arose with a negative sign, the argument would fail. The pairing property is possible precisely because the Petersen graph P_{10} has no 3-cycles. If one added an additional edge to P_{10} , it would have caused a 3-cycle to be created, so the argument would have then failed.) We now return to

$$Z := 12s_5 - 5s_2s_3 + 5s_3\sqrt{4s_4 - s_2^2}.$$

If $12s_5 < 5s_2s_3$ then (2.2) holds and, using Step 2, we obtain

$$\begin{aligned} Z &\geq 30 \sum_{i=1}^{10} \sum_{j=1, i \neq j}^{10} x_i y_j - 30 \sum_{i=1}^{10} x_i y_i + 5s_3[2(x_1 - (x_2 + \cdots + x_{10}))] \\ &= 30(x_1 + x_2 + \cdots + x_{10})(y_1 + y_2 + \cdots + y_{10}) \\ &\quad - 60(x_1 y_1 + x_2 y_2 + \cdots + x_{10} y_{10}) \\ &\quad + 30(x_1 - x_2 - \cdots - x_{10})(y_1 + y_2 + \cdots + y_{10}) \\ &= 60x_1(y_1 + y_2 + \cdots + y_{10}) - 60(x_1 y_1 + x_2 y_2 + \cdots + x_{10} y_{10}) \\ &\geq 0, \end{aligned}$$

since $x_1 y_i \geq x_i y_i$ for $i = 1, 2, \dots, 10$. So $Z \geq 0$ for all trace zero nonnegative 5×5 matrices. This concludes the proof of the theorem.

3. Main results

The following theorem completely solves the NIEP for $n = 5$ when the sum of the five eigenvalues is zero.

Theorem 3.1 (Main theorem). *Let $\lambda_1, \lambda_2, \lambda_3, \lambda_4$ and λ_5 be complex numbers and let $s_k = \lambda_1^k + \lambda_2^k + \lambda_3^k + \lambda_4^k + \lambda_5^k$ for $k = 1, 2, 3, \dots$. Assume $s_1 = 0$. Then $(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5)$ is the spectrum of a nonnegative 5×5 matrix if and only if the following conditions hold:*

1. $s_k \geq 0$ for $k = 2, 3, 4, 5$
2. $4s_4 \geq s_2^2$, and
3. $12s_5 - 5s_2s_3 + 5s_3\sqrt{4s_4 - s_2^2} \geq 0$.

Proof. By [2] and Theorem 2.1 above, the conditions are necessary. So, suppose that the conditions hold. Consider

$$A := \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ q & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ v & u & t & s & 0 \end{pmatrix}.$$

The characteristic polynomial of A is

$$x^5 - (q + s)x^3 - tx^2 + (qs - u)x + qt - v.$$

The Newton identities state that

$$ka_{n-k} + s_1a_{n-k+1} + s_2a_{n-k+2} + \cdots + s_ka_n = 0$$

for $k = 1, 2, \dots, n$ where $p(x) = a_nx^n + a_{n-1}x^{n-1} + \cdots + a_1x + a_0$ with $a_n = 1$ is a monic polynomial with zeroes $\lambda_1, \lambda_2, \dots, \lambda_n$ including multiplicities and $s_k = \lambda_1^k + \lambda_2^k + \cdots + \lambda_n^k$. Applying these identities for $k = 1, 2, \dots, 5$ to the characteristic polynomial of A , we require

$$q + s = s_2/2. \quad (3.1)$$

$$t = s_3/3. \quad (3.2)$$

$$qs - u = \left(\frac{s_2^2}{2} - s_4 \right) / 4. \quad (3.3)$$

$$qt - v = \frac{s_2s_3}{6} - \frac{s_5}{5}. \quad (3.4)$$

Now, Eqs. (3.1) and (3.2) reduce to the inequalities

$$\begin{aligned} qs &\geq \frac{s_2^2}{8} - \frac{s_4}{4}, \\ q \frac{s_3}{3} &\geq \frac{s_2s_3}{6} - \frac{s_5}{5} \\ &= \frac{5s_2s_3 - 6s_5}{30}. \end{aligned}$$

Take $q = s = s_2/4$. Then the inequalities hold if

$$\frac{s_2s_3}{12} \geq \frac{5s_2s_3 - 6s_5}{30},$$

that is, if $12s_5 \geq 5s_2s_3$. Then taking $\mu = (4s_4 - s_2^2)/16$, $v = (12s_5 - 5s_2s_3)/60$, the resulting matrix

$$A := \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ \frac{s_2}{4} & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \frac{12s_5 - 5s_2s_3}{60} & \frac{4s_4 - s_2^2}{16} & \frac{s_3}{3} & \frac{s_2}{4} & 0 \end{pmatrix}$$

is nonnegative and has the given spectrum.

In the general case, we require

$$q + s = s_2/2,$$

$$qs \geq \frac{s_2^2}{8} - \frac{s_4}{4},$$

$$qs_3 \geq \frac{5s_2s_3 - 6s_5}{10}.$$

Write $s = s_2/2 - q$ and substitute into the second inequality to get

$$\frac{s_2q}{2} - q^2 \geq \frac{s_2^2}{8} - \frac{s_4}{4}.$$

Completing the square we get

$$(q - \frac{s_2}{4})^2 \leq \frac{s_4}{4} - \frac{s_2^2}{16} = \frac{1}{16}(4s_4 - s_2^2).$$

The choice

$$q = \frac{s_2}{4} + \frac{1}{4}\sqrt{4s_4 - s_2^2}$$

gives

$$qs_3 = \frac{1}{4}(s_2s_3 + s_3\sqrt{4s_4 - s_2^2}).$$

Hence,

$$qs_3 \geq \frac{1}{10}(5s_2s_3 - 6s_5),$$

since $12s_5 - 5s_2s_3 + 5s_3\sqrt{4s_4 - s_2^2} \geq 0$. This choice of q thus satisfies the conditions provided $s \geq 0$. That is, we require

$$\frac{1}{4}\sqrt{4s_4 - s_2^2} \leq \frac{s_2}{4}$$

(since $q + s = s_2/2$) and thus that $2s_4 \leq s_2^2$. In this case

$$A = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ \frac{s_2}{4} + \frac{1}{4}\sqrt{4s_4 - s_2^2} & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \frac{12s_5 - 5s_2s_3 + 5s_3\sqrt{4s_4 - s_2^2}}{60} & 0 & \frac{s_3}{3} & \frac{s_2}{4} - \frac{1}{4}\sqrt{4s_4 - s_2^2} & 0 \end{pmatrix}$$

is a nonnegative matrix with the given spectrum. Suppose finally that $2s_4 > s_2^2$. Consider

$$A := \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ \frac{s_2}{2} & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ \frac{s_5}{5} & \frac{2s_4 - s_2^2}{8} & \frac{s_3}{3} & 0 & 0 \end{pmatrix}.$$

Then A is nonnegative and has the desired spectrum. This concludes the proof of the theorem.

Acknowledgements

We wish to thank Raphi Loewy for his helpful comments and especially for carefully reading an earlier draft of this paper.

References

- [1] A. Berman, R. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*, Academic Press, New York, 1979.
- [2] T.J. Laffey, E. Meehan, A refinement of an inequality of Johnson, Loewy and London on nonnegative matrices and some applications, *ELA* 3 (1998) 119–128.
- [3] R. Loewy, D. London, A note on the inverse eigenvalue problems for nonnegative matrices, *Linear and Multilinear Algebra* 6 (1978) 83–90.
- [4] H. Mine, *Nonnegative Matrices*, Wiley, New York, 1988.
- [5] R. Reams, *Topics in Matrix Theory*, Thesis presented for the degree of Ph.D., National University of Ireland, Dublin, 1994.
- [6] R. Reams, An inequality for nonnegative matrices and the inverse eigenvalue problem, *Linear and Multilinear Algebra* 41 (1996) 367–375.